



API-MO.BRU


# CAGE VAR-CONTROL



## ENGLISH INSTRUCTIONS FOR USE 1) Insertion of the cage

Put the cage on the foundation on time or insert it on a comb at the centre of the beehive, cutting the part of comb that is sufficient to contain the small cage without cutting the threads of the framework (see figure).

Keep the comb always at the centre of the bees colony. This small cage can remain inside the beehive permanently and it can be also used in the swarming period: putting the queen bee in the cage just before the bees breed royal cells and keeping her in this state for ten days or more, the swarming can be reduced.



## 2) Insertion of the queen

When you insert the queen in the small cage you must be careful with smoke, the queen must not be irritated neither before nor after, especially when she is inside the small cage since she has no way out.

To find the queens easier we advise you to mark them before the putting in the cage, in this way you can understand, at the moment of the putting in cage, if the queen has recently been substituted, such a thing is a clear sign that the beehive is in crisis, since there is the varroa, such a thing happens very frequently.

## 3) Restriction of brood period

To make the putting in the cage at the appropriate moment see "Methods for the putting in cage of the queen for restriction of brood".

To set free the queen bee it is sufficient to open the small door of the small cage and then re-insert the honeycomb to its place.

Attention: do not let the queen bee go out with smoke or other, she will go out spontaneously inside the beehive.

Close the small door after some days to avoid that the bees build inside the small cage. Such a thing happens, if the operation is carried out in spring-summer, on the contrary, if this takes place in autumn, the re-closing of the small door can take place next spring.

### Instructions

Be sure that the whole beehive is normal at the moment of the putting in the cage: if in some beehives the queen has been recently replaced (the marked one is not found and there is a young one) be sure that the new queen lays before being put in the cage. It can also happen that the queen is near to the replacement, because she is compromised by a great load of varroa. This can be understood from the presence of royal cells. If this will happen to put in cage the old queen in any case, without damaging the royal cells. After, the bees will choose which is the queen to keep. Finally, before the chemical treatment, if the new queen has laid a little brood, this must be removed. As concerns this, it is useful to remind that from the arrival of varroa, in August/September the beehives are already compromised, if during the previous year have not been treated well.




API-MO.BRU

## OPENING AND CLOSING OF THE SMALL CAGE


- To close the small cage it is sufficient to make a light pressure on the indicated points,

- To open it, one must work on the same points using the levers.




## INSERTION OF THE VAR-CONTROL CAGE ON THE FOUNDATION


1) weld the foundation to the frame;



2) supply yourself with a 2 cm thickness and 26 cm x 41 cm dimension plank;



3) lay the frame with the welded foundation over the plank then with a blade knife cut into the rectangle that will contain the cage.



4) It is also useful a 2cm thickness and 5.5 x 9cm dimension plank, that is to be laid over the foundation in order to cut a rectangle.




At the moment of the putting in cage it is better if the beehives have already the cages inserted inside, so that must introduced only the queen to start the restriction of brood. In this way times are speeded up.

It is better if the beekeeper prepares in the laboratory, the foundations or the brood combs built with the cages already inserted and that he introduces them in the beehive in advance.

NOTE: while inserting into the beehive a foundation with the already inserted cage, you must consider that the foundation must be worked by the bees. The right period is Spring/Summer. In the period Summer/Autumn the cage must be inserted on the cage built on the brood comb.






WOOD CUBE mm 25 x 43 x 68

It is useful to keep bees from building honeycomb inside the cage .

Keep the front wall closed and the little door is, then draw the hook position on the piece of wood.



Remove some material from the cube wood to give space to the door hook.

N.B to move the cube of wood from the cage pull on the points showed in the picture. DO NOT try to move the cube of wood away from the cage using any tools trough the holes.

API-MO.BRU di Mozzato Bruno Via Palazzon, 48 35010 Campodoro (PD)  
Tel .049/9065542 [www.apimobru.com](http://www.apimobru.com) [info@apimobru.com](mailto:info@apimobru.com)



Beehive Nr	Days following the treatment											
	1	2	3	4	5	6	7	8	9	10	11	12
3	1500	414	75	29	13	5	4	0	1	2	0	0
5	1670	247	72	34	17	10	4	2	1	1	2	0
6	1800	600	66	50	57	31	17	11	9	10	2	2
7	1400	350	38	24	22	9	6	0	2	3	2	0
8	1800	600	47	34	27	12	8	9	3	2	2	0
10	1100	/	/	/	332	64	24	12	16	11	3	2

*Table 1:* number of varroas fallen the days following the treatment with dripped oxalic acid (100 gr of oxalic acid, a Kg of sugar melted in a litre of water).



OTHER DOCUMENTS TO FOLLOW



## METHOD TO CONTROL VARROA ( RESUME OF OUR EXPERIENCE)

### YEARS from 2000 to 2006

- Colonies in brood less state once only, obtained by trapping the queen for 4 weeks
- Queens trapped in cage during the first 10 days of September
- Treatment performed after 24 days, in the first days of October
- Queens released on the 28<sup>th</sup> day.

Number of varroa mites fallen in the first day: from 2000\* to 3000\*.

**2007** (a particular year, warm winter, anticipated spring, 2 treatments, in brood less state, performed)

1<sup>st</sup> brood less state: queen trapped for 24 days:

- Queens trapped in cage on July the 8<sup>th</sup> **leaving the honey super in place to let the bees store honey and removing the queen excluder, in this way the queen will not feel oppressed during the hottest hours.**
- Honey super removed before the treatment.
- All queens were released and the treatments were performed on the same day: the 1<sup>st</sup> of August, after 24 days of trapping.

Treatment performed with drops of oxalic acid (recipe: 100gr oss./1000 gr sugar/one litre of water)

5 cc each favus with bees, dripped for more than half on the port honeycomb slats and the rest on bees (oxalic distributed trickling sideways compared to honeycombs)

Beehive Numbers of varroa mites fallen from the first day to the twelfth day

N°	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°
3	1500*	414	75	29	13	5	4	0	1	2	0	0
5	1670	247	72	34	17	10	4	2	1	1	2	0
6	1800*	600*	66	50	57	31	17	11	9	10	2	2
7	1400*	350*	38	24	22	9	6	0	2	3	2	0
8	1800*	600*	47	34	27	12	8	9	3	2	2	0
10	1100*	----	----	----	332	64	24	12	16	11	3	2

2<sup>nd</sup> brood less state obtained by trapping the queen in cage four weeks:

- Queens trapped on October the 1<sup>st</sup>
- Treatment on October the 25<sup>th</sup>
- Queens released on October the 29<sup>th</sup>

Treatment with oxalic acid dripped (solution 100 gr oss. / 1000 gr sugar / 1 liter of water),

5 cc per honeycomb covered with bees, dripped for more than a half on the port honeycomb slats and the rest on bees.

(oxalic distributed trickling sideways compared to honeycombs)

Beehive Numbers of varroa mites fallen from the first day to the eleventh day

N°	1°	2°	3°	4°	5°	6°	7°	8°	9°	10°	11°	12°
3	207	137	33	21	1	----	----	2	----	2	0	----
5	775	278	20	7	1	----	----	0	----	1	0	----
6	337	292	44	16	2	----	----	2	----	0	0	----
7	260	308	146	76	10	----	----	4	----	1	1	----
8	116	186	96	62	16	----	----	12	----	5	2	----
10	359	356	104	55	11	----	----	14	----	1	4	----

\* approximate numbers

---- days in which no count was performed

Taking into consideration the hive # 5: 1670 varroa mites fell on the 1<sup>st</sup> day, a total of 2060 varroa mites fell until the 12<sup>th</sup> day (considering that varroa doubles from month to month) and considering that for about 2/3 of July varroa could not multiply due to the brood less state, we can foresee the number of varroa mites possibly reached on August the 1<sup>st</sup>:  $(2060: 3 \times 2) = 1373$ , we add  $2060 + 1373 = 3433$  varroa mites (number of varroa mites beyond which we would have the crisis of the colony the next month). Thanks to the two brood less states performed, one short in July and a longer one at the end of the season in October, I was able to sanitize my colonies.

NOTE: making a comparison of the years 2000 to 2006 with 2007, you notice that in 2007 varroa reached the same numbers 2 months before, if I waited one month more in intervening, the colonies could have been compromised .

The method adopted in 2007 and in the subsequent years, 2022 included, considers two periods with queen bee in the cage: one in summer (24 days long, starting on the days between July 10 and July 15) and one in autumn (28 days long, starting on the days between October 5 and October 15). The results are excellent: varroa mite hovered at low numbers throughout the year.

### **When did I trapped queens?**

- One time In July, immediately after the honey storing, when the bees reached the maximum growth. From this point on, because of the heat and the lack of honey storing activities, the bees decreased the brood rearing. So, that is the most appropriate time to intervene. Thanks to the total absence of brood I apply a treatment against varroa of proven efficacy in the first days of August. You should not wait until mid-August to perform the caging, varroa may have reached high numbers and as a result brood can be heavily infested and will not give healthy bees. As a consequence the colony will collapse, when the bees born among May, June and July, will die at the end of their life cycle;

- **In the fall (October), a month before the Winter cluster, another queen entrap:** I make all hives, at the same time and as soon as possible, without brood and capable of receiving the treatment against varroa very soon, in the first days of November, not in December or January, when the varroa will have done irreparable damage.

Bees from November onwards are almost still as assets and also considering breeding brood, brood rearing can occur if the temperatures are good but it is always a small amount, that won't create the future colony. On the other hand, this little brood will be full of varroa mites, newborn bees will be debilitated and unable to flicker alone. The bees have made a useless work of rearing brood and also it will be difficult to clean them from varroa after the season.

Conclusion: contain varroa mite at low numbers to maintain healthy colonies.

To sum up, targeted treatments, in total absence of brood, thanks to queen caging is the ideal condition to reduce varroa:

- One treatment in summer to hover the number of varroa, in order to avoid the damage that high numbers of them will cause. **Beware:** we are still in active season for bees so, reinfestation can happen;

- An autumn cleaning at the end of the season, easy to apply because the hive is no longer crowded with bees, perform it when bees are retreating home and there is no longer the risk of re-infestation among apiaries because of drones or looting, I consider it essential for a good awakening in the next year, especially in areas where bees go on with brood rearing and never stop. The total absence of brood planned thanks to queen caging, is helpful in breaking down varroa, in order to keep bees healthy. Moreover, it allows you to avoid the repetition of the treatments several times that weaken the colonies.

Thinking about what dates it is better to perform the queen entrap and subsequent treatment, every beekeeper will have their own opinion, in accordance with the climate in their area.

The result of this method, confirmed also by beekeepers that have adopted it are:

- Colonies full of bees each year;
- Healthy bees in the fall that can easily pass through the winter;
- Excellent recovery in spring;
- Excellent honey production;